

Diversity of Individual Mobility Patterns

Xiao-Yong Yan^{1,2,3,4}, Xiao-Pu Han^{3,5}, Bing-Hong Wang³, and Tao Zhou^{1,3*}

¹*Web Sciences Center, University of Electronic Science and Technology of China, Chengdu 611731, P.R. China*

²*Department of Transportation Engineering, Shijiazhuang Tiedao University, Shijiazhuang 050043, P.R. China*

³*Department of Modern Physics, University of Science and Technology of China, Hefei 230026, P.R. China*

⁴*Department of Systems Science, Beijing Normal University, Beijing 100875, P.R. China*

⁵*Institute of Information Economy and Alibaba Business College,
Hangzhou Normal University, Hangzhou 310036, P.R. China*

Uncovering human mobility patterns is of fundamental importance to the understanding of epidemic spreading, urban transportation and other socioeconomic dynamics embodying spatiality and human travel. The observed scaling laws for aggregated data require a theoretical explanation of their underlying mechanism. According to the direct travel diaries of volunteers, we show the absence of scaling properties in the displacement distribution at the individual level, which unfortunately provides a complete contrast to most inferences and assumptions in the literature. The aggregated displacement distribution follows a power law with an exponential cutoff, which is analytically explained by the mixture nature of human travel under Maxwell-Boltzmann statistics. Our analysis provides an alternative way to bridge diverse patterns at the individual level and scaling laws at the population level.

I. INTRODUCTION

Positioning systems in mobile phones and vehicles and Wi-Fi devices in laptop computers and personal digital assistants have made quantitative analyses of human mobility patterns possible [1–3]. These analyses have a significant potential to reveal novel statistical regularities of human behavior, refine our understanding of the socioeconomic dynamics embodying spatiality and human mobility [4, 5], and eventually contribute to controlling disease [6–9], designing transportation systems [10], locating facilities [11], providing location-based services [12, 13], and so on.

Aggregated data from bank notes [14], mobile phones [1] and onboard GPS measurements [2] showed that the displacement distribution of human mobility, for both long-range travel and daily movements, approximately follows a power law. The scaling laws in long-range travel may result from the hierarchical organization of transportation systems [15], while the scaling laws in daily movements have recently been explained by the *exploration and preferential return* mechanism [16]. This model [16] suggested that the displacement distribution at the individual level is power-law, which could also be considered to be a straightforward inference from the results of aggregated data.

Thus far, we still lack solid results about human mobility patterns at the individual level. Inferring individual features from the aggregated data is very risky because the scaling law for the population could be a mixture of many individuals with different statistics [17]. In addition, the aforementioned data are not sufficient to draw conclusions at the individual level. First, data such as GPS records from taxis and the trajectories of bank

notes consist of many individual movements, but these individuals are not easy to be distinguished from each other. Second, data such as GPS records from mobile phones and the trajectories of bank notes could not accurately capture purposeful travels with explicit origins and destinations. In fact, the displacement between two activations of a mobile phone may be just a tiny portion of a purposeful trip or a combination of several sequential trips, while the displacement between two registrations of a bank note could be the result of a number of sequential trips made by different people.

Instead of using proxy data, we analyze the travel diaries of hundreds of volunteers. Though the data set is small, it contains personal profiles and explicit positions of origins and destinations, allowing quantitative and authentic analyses at the individual level. In contrast to the scaling laws in aggregated data, people show diverse mobility patterns, and few of them display the scaling property. In fact, the trajectories of students and employees are dominated by trips connecting homes with schools and workplaces, respectively, while trips are distributed more homogeneously among different locations for others such as retirees, homemakers, unemployed people, and so on. The aggregated displacement distribution follows a power law with an exponential cutoff, which can be analytically explained by the mixed nature of human travel under Maxwell-Boltzmann statistics. In addition, this theory predicts that the displacements using a single means of transportation will follow an exponential distribution, which is also supported by the empirical data on taxi trips and air flights.

* zhutou@ustc.edu

II. RESULTS

A. Individual mobility patterns

Our analysis of human mobility is based on a data set of 230 volunteers' six-week travel diaries in Frauenfeld, Switzerland [18]. This data set contains the volunteers' personal information, including age, job and sex, and 36761 trip records. Each record includes the geographic positions (longitude and latitude) of a trip's origin and destination. By calculating the spherical distance between two trip endpoints from their longitudes and latitudes, we can obtain the length of each trip.

We first measure the individual displacement distributions from the data set. Figs. 1(a)-1(c) show three typical individuals' displacement distributions (**Table S1** presents all volunteers' displacement distributions), from which we cannot find any universal scaling properties. Indeed, when we use the *Kolmogorov-Smirnov* test [19] to test whether the distributions fit the power law, we find that 87.8% of the individuals cannot pass the test (statistical validation results are listed in **Table S2**). This result strongly suggests the absence of scaling laws in human travel at the individual level.

To reveal the underlying structure of individual trips, we assign to each individual a mobility network, in which the nodes denote locations visited by individuals, edges represent the trips between nodes and edge weight is defined as the number of corresponding trips [20]. Figs. 1(d)-1(f) show three typical individuals' mobility networks (all networks are presented in **Table S1**). As shown in Fig. 1 and **Table S1**, for most students and employees, their edge weights are highly heterogeneous. For each individual, we call the trip corresponding to the edge with the largest weight the *dominate trip* and define the domination ratio d as the ratio of the weight of the dominate trip to the total weight. Fig. 2 reports the distribution of domination ratios for different groups of individuals, from which we can see that the student group has the largest d on average and that the employees' average domination ratio is smaller than that of the students but larger than that of the other group.

The difference of d results from the fact that students and employees frequently travel between homes and schools/workplaces in working days but retirees or homemakers do not have to do so. The peak values in the displacement distributions of students and employees are thus usually determined by the lengths of their dominant trips. Because the lengths of dominant trips are not necessarily small, the displacement distribution for an individual is usually not right-skewed and is far different from a power law. In addition, the significant role of the dominant trip indicates that an individual's traveling process in general cannot be characterized by the Lévy flight or truncated Lévy flight.




FIG. 1. Individual mobility patterns. **a-c**. Displacement distributions for three typical individuals (**a** - a student, **b** - an employee, **c** - a retiree), where the peak values for the student and the employee result from the trips between two most frequently visited locations. **d-f**. Mobility networks for the three individuals, where the area of a node is proportional to its number of visits and the width of an edge is proportional to its weight.




FIG. 2. Distribution of the domination ratios. **a**. Population. **b**. Student group. **c**. Employee group. **d**. Others. S is the number of group members, and $\bar{d}$ is the average domination ratio.




FIG. 3. Displacement distribution $P(r)$ of the aggregated data. The solid line indicates a power law with an exponential cutoff.

B. Scaling property in aggregated data

The aggregated displacement distribution of individuals (see Fig. 3) is well approximated by a power law with an exponential cutoff $P(r) \propto r^{-1.05} \exp(-r/50)$, which is similar to those observed for bank notes [14] and mobile phone users [1]. As shown above, this scaling property is not a simple combination of many analogous individuals. We assume that the total travel cost is C , the number of trips with cost c_i is n_i , and the same-cost trips are indistinguishable, though they could be made by different individuals. Recalling *Maxwell-Boltzmann* statistics, the number of microstates is $\Omega = N! (\prod_i n_i!)^{-1}$ where N is the total number of trips. According to the *maximum entropy principle* [21], the maximization of Ω (under two constraints, $\sum n_i = N$ and $\sum n_i c_i = C$) leads to the solution $n_i \propto \exp(-c_i/\bar{c})$, where $\bar{c} = C/N$ is the average travel cost. Denote the density of trips with cost c by $P(c)$, then $P(c) \propto \exp(-c/\bar{c})$.

The travel cost is commonly approximated as the weighted sum $c \approx \eta t + \mu m$, where η and μ are two coefficients, and t and m are the costs involving time and money, respectively. Previous empirical studies have suggested that the monetary cost is approximately proportional to the travel distance as $m \approx \nu r$ [22], while the travel time approximately obeys a logarithmic form, $t \approx \phi \ln r + \psi$ [23, 24], where ν , ϕ and ψ are coefficients. The logarithmic relation results from the mixture of modes of transportation [25]. Apparently, people move faster when traveling longer distances: we walk from the classroom to the restaurant but take an airplane from the US to China. Integrating the aforementioned terms, we obtain the displacement distribution $P(r) \propto r^{-\beta} \exp(-r/\kappa)$, where $\beta = \eta\phi/\bar{c}$ and $\kappa = \bar{c}/\mu\nu$.

A direct corollary of Maxwell-Boltzmann statistics is that the displacement distribution should follow an exponential form if it only accounts for trips from a sin-

gle mode of transportation because in that case, $c \propto r$. This corollary has found strongly supportive evidence from a number of empirical studies on disparate systems [26–30]. **Figure S1** reports empirical distributions for taxi trajectories in Beijing [27], car trajectories in New York (downloaded from *nhts.ornl.gov*), bus trips in Shijiazhuang (collected by the authors) and air flights in the US [28]. All distributions can be well characterized by exponential-like functions.

III. DISCUSSION

The general lessons that we learned from the present analysis could be used to refine our knowledge of human mobility patterns. First, the displacement distributions for aggregated data usually display power-law decay with an exponential cutoff. Meanwhile, there are examples ranging from taxi trips to air flights in which the displacement distributions are exponential. In these examples, every displacement distribution is generated by trips involving a single mode of transportation, which corresponds to a linear relation between the travel cost and distance and eventually results in an exponential displacement distribution according to Maxwell-Boltzmann statistics. The present results suggest that the form (power law or exponential or other) of deterrence function in the gravity law for human travel [31] may be sensitive to the modes of transportation under consideration.

This study warns researchers of the risk of inferring individual behavioral patterns from aggregated statistics. Analogously, the temporal burstiness of human activities is widely observed, and the researchers are aware of the fact that the aggregated scaling laws could either be a combination of a number of individuals, each of whom displays scaling laws similar to the population [32], or the result of a mixture of diverse individuals, most of whom exhibit far different statistical patterns than the population [33–35]. In comparison, such issues are less investigated for spatial burstiness. In particular, experimental analyses on individuals has rarely been reported. Determining whether the displacement distribution of an individual follows a power-law distribution will require further data and analysis.

Many known mechanisms underlie the scaling laws of complex systems [36–38], including rich get richer [39, 40], good get richer [41, 42], merging and regeneration [43], optimization [44, 45], Hamiltonian dynamics [46], stability constraints [47], and so on. The individual mobility model by Song *et al.* [16] is a typical example embodying the rich get richer mechanism. The maximum entropy theory under Maxwell-Boltzmann statistics gives an unpretentious yet reasonable explanation for the emergent scaling from diverse individuals. In summary, this work is complementary to known results on human mobility patterns and provides insights into how to bridge the gap between diverse individual statistics and aggregated regular patterns.

-
- [1] González, M. C., Hidalgo, C. A. & Barabási, A. L. Understanding individual human mobility patterns. *Nature* **453**, 779-782 (2008).
 - [2] Jiang, B., Yin, J. & Zhao, S. Characterizing the human mobility pattern in a large street network. *Phys. Rev. E* **80**, 021136 (2009).
 - [3] Yoon, J., Noble, B. D., Liu, M. & Kim, M. Building realistic mobility models from coarse-grained traces. in: *Proc. of the ACM MobiSys'06*, (Uppsala, Sweden), pp 177-190 (2006).
 - [4] Vespignani, A. Predicting the behavior of techno-social systems. *Science* **325**, 425-428 (2009).
 - [5] Barthélemy, M. Spatial networks. *Phys. Rep.* **499**, 1-101 (2011).
 - [6] Balcan, D. & Vespignani, A. Phase transitions in contagion processes mediated by recurrent mobility patterns. *Nat. Phys.* **7**, 581-586 (2011).
 - [7] Belik, V., Geisel, T. & Brockmann, D. Natural human mobility patterns and spatial spread of infectious diseases. *Phys. Rev. X* **1**, 011001 (2011).
 - [8] Ni, S. & Weng, W. Impact of travel patterns on epidemic dynamics in heterogeneous spatial metapopulation networks. *Phys. Rev. E* **79**, 016111 (2009).
 - [9] Zhao, Z.-D., Liu, Y. & Tang, M. Epidemic variability in hierarchical geographical networks with human activity patterns. *Chaos* **22**, 023150 (2012).
 - [10] Horner, M. W. & O'Kelly, M. E. S. Embedding economies of scale concepts for hub networks design. *J. Transp. Geogr.* **9**, 255-265 (2001).
 - [11] Um, J., Son, S.-W., Lee, S.-I., Jeong, W. & Kim, B. J. Scaling laws between population and facility densities. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 14236-14240 (2009).
 - [12] Clements, M., Serdyukov, P., de Vries A. P. & Reinders, M. J. T. Personalised travel recommendation based on location co-occurrence. arXiv:1106.5213
 - [13] Scellato, S., Noulas, A. & Mascolo C. Exploiting place features in link prediction on location-based social networks. in: *Proc. of the ACM KDD'11*, (New York, USA), pp 1046-1054 (2011).
 - [14] Brockmann, D., Hufnagel, L. & Geisel, T. The scaling laws of human travel. *Nature* **439**, 462-465 (2006).
 - [15] Han, X.-P., Hao, Q., Wang, B.-H. & Zhou, T. Origin of the scaling law in human mobility: hierarchy of traffic systems. *Phys. Rev. E* **83**, 036117 (2011).
 - [16] Song, C., Koren, T., Wang, P. & Barabási, A. L. Modelling the scaling properties of human mobility. *Nat. Phys.* **6**, 818-823 (2010).
 - [17] Petrovskii S., Mashanova A. & Jansen, V. A. A. Variation in individual walking behavior creates the impression of a Lévy flight. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 8704-8707 (2011).
 - [18] Chalasani, V. S., Engebretsen, Ø. Denstadli, J. M. & Axhausen, K.W. Precision of geocoded locations and network distance estimates. *J. Transport. Stat.* **8**, 1-15 (2005).
 - [19] Clauset, A., Shalizi, C. R. & Newman, M. E. J. Power-law distributions in empirical data. *SIAM Rev.* **51**, 661-703 (2009).
 - [20] Song, C., Qu, Z., Blumm, N. & Barabási, A. L. Limits of predictability in human mobility. *Science* **327**, 1018-1021 (2010).
 - [21] Balescu, R. Equilibrium and nonequilibrium statistical mechanics. (New York: John Wiley), (1975).
 - [22] Willumsen, L. G. Travel networks. in *Handbook of Transport Modelling*, eds Hensher, D. A. & Button, K. J. (New York: Pergamon), pp 165-180 (2000).
 - [23] Rietveld, P., Zwart, B., van Wee, B. & van den Hoorn, T. On the relationship between travel time and travel distance of commuters. *Ann. Reg. Sci.* **33**, 269-287 (1999).
 - [24] Li, S., Wang, H. & Wang, Z. A study on tour time planning of domestic sightseeing travel itineraries. *Hum. Geogr.* **20**, 51-56 (2005).
 - [25] Oosterhaven, J. A. & Rietveld, P. Transport costs, location and the economy. in: *Location and Competition*, eds Brakman, S. & Garretsen, H. (New York: Routledge), pp 32-60 (2005).
 - [26] Roth, C., Kang, S. M., Batty, M. & Barthélemy, M. Structure of urban movements: polycentric activity and entangled hierarchical flows. *PLoS ONE* **6**, e15923 (2011).
 - [27] Liang, X., Zheng, X., Lü, W., Zhu, T. & Xu, K. The scaling of human mobility by taxis is exponential. *Physica A* **391**, 2135-2144 (2012).
 - [28] Jiang, B. & Jia, T. Exploring human mobility patterns based on location information of US flights. arXiv:1104.4578v2
 - [29] Gallotti, R., Bazzani, A. & Rambaldi, S. Towards a Statistical physics of human mobility. *Int. J. Mod. Phys. C* **23**, 1250061 (2012).
 - [30] Peng, C., Jin, X., Wong, K. C., Shi, M. & Liò, P. Collective human mobility pattern from taxi trips in urban area. *PLoS ONE* **7**, e34487 (2012).
 - [31] Simini, F., González, M. C., Maritan, A. & Barabási, A.-L. A universal model for mobility and migration patterns. *Nature* **484**, 96-100 (2012).
 - [32] Barabási, A. L. The origin of bursts and heavy tails in human dynamics. *Nature* **435**, 207-211 (2005).
 - [33] Malmgren, R. D., Stouffer, D. B., Motter, A. E., & Amaral L. A. N. A Poissonian explanation for heavy tails in e-mail communication. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 18153-18158 (2008).
 - [34] Hidalgo, C. A. Conditions for the emergence of scaling in the inter-event time of uncorrelated and seasonal systems. *Physica A* **369**, 877-883 (2006).
 - [35] Wu, Y., Zhou, C., Xiao, J., Kurths, J. & Schellnhuber, H. J. Evidence for a bimodal distribution in human communication. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 18803-18808 (2010).
 - [36] Mitzenmacher, M. A brief history of generative models for power law and lognormal distributions. *Internet Math.* **1**, 226-251 (2004).
 - [37] Newman, M. E. J. Power laws, Pareto distributions and Zipf's law. *Contemp. Phys.* **46**, 323-351 (2005).
 - [38] Simkin, M. V. & Roychowdhury, V. P. Re-inventing Willis. *Phys. Rep.* **502**, 1-35 (2011).
 - [39] Simon, H. A. On a class of skew distribution functions. *Biometrika* **42**, 425-440 (1955).
 - [40] Barabási, A. L. & Albert, R. Emergence of scaling in random networks. *Science* **286**, 509-512 (1999).
 - [41] Garlaschelli, D., Capocci, A. & Caldarelli, G. Self-organized network evolution coupled to extremal dynamics. *Nat. Phys.* **3**, 813-817 (2007).

- [42] Lü, L., Zhang, Z.-K. & Zhou, T. Zipf's Law Leads to Heaps' Law: analyzing their relation in finite-size systems. *PLoS ONE* **5**, e14139 (2010).
- [43] Kim, B. J., Trusina, A., Minnhagen, P. & Sneppen, K. Self organized scale-free networks from merging and regeneration. *Eur. Phys. J. B* **43**, 369-372 (2005).
- [44] Valverde, S., Cancho, F. & Solé, R. V. Scale-free networks from optimal design. *Eur. Phys. Lett.* **43**, 369-372 (2002).
- [45] Bartumeus, F., Da Luz, M. G. E., Viswanathan, G. M. & Catalan, J. Animal search strategies: a quantitative random-walk analysis. *Ecology* **86**, 3078-3087 (2005).
- [46] Baiesi, M. & Manna, S. Scale-free networks from a Hamiltonian dynamics. *Phys. Rev. E* **68**, 047103 (2003).
- [47] Perotti, J. I., Billoni, O. V., Tamarit, F. A., Chialvo, D. R. & Cannas, S. A. Emergent self-organized complex network topology out of stability constraints. *Phys. Rev. Lett.* **103**, 108701 (2009).